



itDose

Taking You Beyond The Limits

Visit ID : ITD766
UHID/MR No : ITD.0000000686
Patient Name : Mr.VDUMMY
Age/Gender : 22 Y O M 1 D /M
Ref Doctor : Dr.SELF
Client Name : STANDARD
Client Add : D 159, 1st Floor, Sector 7, No

Registration : 30/Nov/2022 11:57AM
Collected : 30/Nov/2022 11:58AM
Received : 30/Nov/2022 11:58AM
Reported : 30/Nov/2022 11:58AM
Status : Final Report
Client Code : 78
Barcode No : 10059692

DEPARTMENT OF BIOCHEMISTRY

Test Name	Result	Status	Unit	Bio. Ref. Range	Method
-----------	--------	--------	------	-----------------	--------

SERUM LIPASE

Sample Type : Serum

SERUM LIPASE	58	N	U/L	<=38 U/L	Color
--------------	----	---	-----	----------	-------

COMMENTS:

Lipase measurements are used in the diagnosis and treatment of diseases of the pancreas, such as acute pancreatitis, chronic pancreatitis, and obstruction of the pancreatic duct. In acute pancreatitis serum lipase activity tends to become elevated & remains for about 7 - 10 days. Increased lipase activity rarely lasts longer than 14 days, and prolonged increases suggest a poor prognosis or the presence of a cyst. Serum lipase may also be elevated in patients with chronic pancreatitis, obstruction of the pancreatic duct and non pancreatic conditions including renal diseases, various abdominal diseases such as acute cholecystitis, intestinal obstruction or infarction, duodenal ulcer, and liver disease, as well as alcoholism & diabetic keto-acidosis & in patients who have undergone endoscopic retrograde cholangiopancreatography. Elevation of serum lipase activity in patients with mumps strongly suggests significant pancreatic as well as salivary gland involvement by the disease.

*** End Of Report ***

Dr.VIKAS REDDY
Consultant Pathologist

