



itDose

Taking You Beyond The Limits

<b>Visit ID</b> : ITD773	<b>Registration</b> : 30/Nov/2022 02: 24PM
<b>UHID/MR No</b> : ITD.0000000693	<b>Collected</b> : 30/Nov/2022 02: 27PM
<b>Patient Name</b> : Dr.IMRAN	<b>Received</b> : 30/Nov/2022 02: 29PM
<b>Age/Gender</b> : 33 Y O M O D /M	<b>Reported</b> : 30/Nov/2022 02: 32PM
<b>Ref Doctor</b> : Dr.SELF	<b>Status</b> : Final Report
<b>Client Name</b> : STANDARD	<b>Client Code</b> : 78
<b>Client Add</b> : D 159, 1st Floor, Sector 7, No	<b>Barcode No</b> : 10059697

DEPARTMENT OF BIOCHEMISTRY

Test Name	Result	Status	Unit	Bio. Ref. Range	Method
-----------	--------	--------	------	-----------------	--------

SERUM LIPASE

Sample Type : Serum

SERUM LIPASE	12	N	U/L	<=38 U/L	Color
--------------	----	---	-----	----------	-------

ok

COMMENTS:

Lipase measurements are used in the diagnosis and treatment of diseases of the pancreas, such as acute pancreatitis, chronic pancreatitis, and obstruction of the pancreatic duct. In acute pancreatitis serum lipase activity tends to become elevated & remains for about 7 - 10 days. Increased lipase activity rarely lasts longer than 14 days, and prolonged increases suggest a poor prognosis or the presence of a cyst. Serum lipase may also be elevated in patients with chronic pancreatitis, obstruction of the pancreatic duct and non pancreatic conditions including renal diseases, various abdominal diseases such as acute cholecystitis, intestinal obstruction or infarction, duodenal ulcer, and liver disease, as well as alcoholism & diabetic keto-acidosis & in patients who have undergone endoscopic retrograde cholangiopancreatography. Elevation of serum lipase activity in patients with mumps strongly suggests significant pancreatic as well as salivary gland involvement by the disease.

\*\*\* End Of Report \*\*\*

**Dr.VIKAS REDDY**  
**Consultant Pathologist**

